

The time independent spherically symmetric solution of the Einstein equation revisited

Jean-Pierre Petit, Gilles d'Agostini and Sebastien Michea⁺

Manaty Research group

Key words: Non contractible hypersurface. Throat sphere, space bridge. Spherically symmetric solution.

Abstract: Spherical symmetry does not immediately mean central symmetry. The time independent, spherically symmetric solution of the homogeneous Einstein equation is revisited with coordinates which keep the signature invariant and prevent time and radial coordinate interchange. The associated hypersurface is not contractible and corresponds to a space bridge linking two Minkowski spacetimes through a throat sphere. As the determinant of the metric vanishes on that sphere one gets an orbifold structure. When crossing that sphere the particles experience a PT, mass and energy inversions.

Introduction and main idea of this article.

In 1916 Karl Schwarzschild publishes [1] a solution of the vacuum Einstein equations (without second term) correspond to time translation invariance and spherical symmetry. It is only in 1999 that an English translation of this article will be available [2] thanks to S.Antoci et A.Loinger. Schwarzschild decides to express this solution by using a first set of real variables

$$(a) \quad \{ t, x, y, z \} \in \mathbb{R}^4$$

The solution is then given in the form :

$$(b) \quad ds^2 = F dt^2 - G (dx^2 + dy^2 + dz^2) - H (x dx + y dy + z dz)^2$$

He then introduces an intermediary variable :

$$(c) \quad r = \sqrt{x^2 + y^2 + z^2}$$

Which, given (a) is essentially positive.

He performs a new coordinates change that allows him to simply express the spherical symmetry hypothesis :

$$(d) \quad x = r \sin \theta \cos \varphi \quad y = r \sin \theta \sin \varphi \quad z = r \cos \theta$$

⁺ jean-pierre.petit@manaty.net , gilles.dagostini@manaty.net, sebastien.michea@manaty.net

He then obtains the form :

$$(e) \quad ds^2 = F dt^2 - (G + H r^2) dr^2 - G r^2 (d\theta^2 + \sin^2 \theta d\varphi^2)$$

In order for the metric to be lorentzian at infinity it is necessary that :

$$(f) \quad r \rightarrow \infty \quad \text{implies} \quad F \rightarrow 1, G \rightarrow 1, H \rightarrow 0$$

He introduces next a new change of coordinates :

$$(g) \quad x_1 = \frac{r^3}{3}, \quad x_2 = -\cos\theta, \quad x_3 = \varphi, \quad x_4 = t$$

Note that (a) + (c) imply that $x_1 \geq 0$

In those new coordinates the metric becomes :

$$(h) \quad ds^2 = f_4 dx_4^2 - f_1 dx_1^2 - f_2 \frac{dx_2^2}{1-x_2^2} - f_3 dx_3^2 (1-x_2^2)$$

where $f_1, f_2 = f_3, f_4$ are three functions of x_1 (hence of r) that must satisfy the conditions :

$$\text{- For large } x_1 : \quad f_1 = \frac{1}{r^4} = (3x_1)^{-4/3}, \quad f_2 = f_3 = r^2 = (3x_1)^{2/3}, \quad f_4 = 1$$

$$\text{- The determinant } f_1 f_2 f_3 f_4 = 1$$

- The metric must be a solution of the field equation.

- Except for $x_1 = 0$ the f functions must be continuous.

His computation leads him to :

$$(i) \quad f_1 = \frac{(3x_1 + \alpha^3)^{-4/3}}{1 - \alpha(3x_1 + \alpha^3)^{-1/3}} \quad f_2 = f_3 = (3x_1 + \alpha^3)^{2/3} \quad f_4 = 1 - \alpha(3x_1 + \alpha^3)^{-1/3}$$

α being an integration constant.

Using (g) we can rewrite his solution as :

$$(j) \quad ds^2 = \left[1 - \frac{\alpha}{(r^3 + \alpha^3)^{1/3}} \right] dt^2 - \frac{r^4}{(r^3 + \alpha^3) [(r^3 + \alpha^3)^{1/3} - \alpha]} dr^2 - (r^3 + \alpha^3)^{2/3} (d\theta^2 + \sin^2 \theta d\varphi^2)$$

That we can rewrite :

$$(k) \quad ds^2 = g_{tt} dt^2 - g_{rr} dr^2 - g_{\theta\theta} d\theta^2 - g_{\varphi\varphi} d\varphi^2$$

The coefficients g_{tt} , g_{rr} et $g_{\theta\theta}$ are functions of the only variable r , the quantity $g_{\varphi\varphi}$ being a function of both r and φ . The variable r varies from 0 to infinity while being strictly positive by definition (c).

Let us limit ourself to the case $\alpha > 0$ and let consider a path corresponding to $dt = dr = 0$. It has a perimeter :

$$(l) \quad p = 2\pi(r^3 + \alpha^3)^{1/3}$$

This perimeter has a minimal value $p = 2\pi\alpha$.

The hypersurface solution is therefore non contractible.

The r variable cannot be considered as a « radial distance ». This hypersurface does not have a « centre », the object corresponding to $r = 0$ hence to $x = y = z = 0$ is not a dot but a sphere of radius α .

We will now replicate Scharzschild's calculations by replacing the x, y, z and t variables by Greek characters (ρ is however kept for a later purpose), so that the reader not be tempted to assimilate them to distances (x, y and z) or time (t) and lose sight of what they really are: real numbers.

Revisiting Schwarzschild computation.

Let us consider the zero second member Einstein equation $R_{\mu\nu} = 0$ in time independent and spherically symmetrical conditions. Let ξ_1, ξ_2, ξ_3 stand for rectangular coordinates, and ξ_0 as the time marker, with $(\xi_0, \xi_1, \xi_2, \xi_3) \in \mathbf{R}^4$ which stands real values for all coordinates. In addition we assume that there are no crossed terms in the line element, so that this last can be written :

$$(1) \quad ds^2 = F d\xi_0^2 - G(d\xi_1^2 + d\xi_2^2 + d\xi_3^2) - H(\xi_1 d\xi_1 + \xi_2 d\xi_2 + \xi_3 d\xi_3)^2$$

where F, G, H are functions of $\sqrt{\xi_1^2 + \xi_2^2 + \xi_3^2}$.

At infinity we must have

$$(2) \quad F \text{ and } G \rightarrow 1 \quad H \rightarrow 0$$

Introduce the following coordinate change :

$$(3) \quad \zeta = \sqrt{\xi_1^2 + \xi_2^2 + \xi_3^2}$$

$$(4) \quad \theta = \arccos\left(\frac{\xi_3}{\zeta}\right)$$

$$(5) \quad \varphi = \arccos\left(\frac{\xi_1}{\sqrt{\xi_1^2 + \xi_2^2}}\right)$$

which goes with : $\zeta \in \mathbf{R}_+$ $\theta \in \mathbf{R}$ $\varphi \in \mathbf{R}$ and :

$$\xi_1 = \zeta \sin\theta \cos\varphi, \quad \xi_2 = \zeta \sin\theta \sin\varphi, \quad \xi_3 = \zeta \cos\theta$$

that we will call « pseudo spherical coordinates ». It gives :

$$(6) \quad ds^2 = F d\xi_0^2 - (G + H\zeta^2) d\zeta^2 - G\zeta^2 (d\theta^2 + \sin^2\theta d\varphi^2)$$

Introduce the following additional coordinate change : :

$$(7) \quad \eta_1 = \frac{\zeta^3}{3}, \quad \eta_2 = -\cos\theta, \quad \eta_3 = \varphi$$

Then we have the volume element $\zeta^2 \sin\theta d\zeta d\theta d\varphi = d\xi_1 d\xi_2 d\xi_3$. The new variables are then pseudo polar coordinate with the determinant 1. They have the evident advantage of polar coordinates for the treatment of the problem.

In the new pseudo polar coordinates :

$$(8) \quad ds^2 = F d\xi_0^2 - \left(\frac{G}{\zeta^4} + \frac{H}{\zeta^2}\right) d\eta_1^2 - G\zeta^2 \left[\frac{d\eta_2^2}{1-\eta_2^2} + d\eta_3^2 (1-\eta_2^2) \right]$$

for which we write :

$$(9) \quad ds^2 = f_4 d\xi_0^2 - f_1 d\eta_1^2 - f_2 \frac{d\eta_2^2}{1-\eta_2^2} - f_3 d\eta_3^2 (1-\eta_2^2)$$

Then $f_1, f_2 = f_3, f_4$ are functions of η_1 which have to fullfill the following conditions :

$$1 - \text{For } \xi_1 = \infty : f_1 = \frac{1}{\zeta^4} = (3\eta_1)^{-4/3}, \quad f_2 = f_3 = \zeta^2 = (3\eta_1)^{2/3}, \quad f_4 = 1$$

$$2 - \text{The equation of the determinant : } f_1 \cdot f_2 \cdot f_3 \cdot f_4 = 1$$

3 - The field equations.

4 - Continuity of the f, except for $\eta_1 = 0$

In order to formulate the field equations one must first form the components of the gravitational field corresponding to the line element (9). This happens in the simplest way when one builds the differential equation of the geodesic line by direct execution of the variation, and reads out the components of these. The differential equations of the geodesic line for the line element (9) immediately result from the variation in the form :

(10)

$$0 = f_1 \frac{d^2 \eta_1}{ds^2} + \frac{1}{2} \frac{\partial f_4}{\partial \eta_1} \left(\frac{d\eta_4}{ds} \right)^2 + \frac{1}{2} \frac{\partial f_1}{\partial \eta_1} \left(\frac{d\eta_1}{ds} \right)^2 - \frac{1}{2} \frac{\partial f_2}{\partial \eta_1} \left[\frac{1}{1-\eta_2^2} \left(\frac{d\eta_2}{ds} \right)^2 + (1-\eta_2^2) \left(\frac{d\eta_3}{ds} \right)^2 \right]$$

(11)

$$0 = \frac{f_2}{1-\eta_2^2} \frac{d^2 \eta_2}{ds^2} + \frac{\partial f_2}{\partial \eta_1} \frac{1}{1-\eta_1^2} \frac{d\eta_1}{ds} \frac{d\eta_2}{ds} + \frac{f_2 \eta_2}{(1-\eta_1^2)^2} \left(\frac{d\eta_2}{ds} \right)^2 + f_2 \eta_2 \left(\frac{d\eta_3}{ds} \right)^2$$

(12)

$$0 = f_2 (1-\eta_2^2) \frac{d^2 \eta_3}{ds^2} + \frac{\partial f_2}{\partial \eta_1} (1-\eta_2^2) \frac{d\eta_1}{ds} \frac{d\eta_3}{ds} - 2 f_2 \eta_2 \frac{d\eta_2}{ds} \frac{d\eta_3}{ds}$$

(13)

$$0 = f_4 \frac{d^2 \eta_4}{ds^2} + \frac{\partial f_4}{\partial \eta_1} \frac{d\eta_1}{ds} \frac{d\eta_4}{ds}$$

The comparison with

(14)

$$\frac{d^2 \eta_\alpha}{ds^2} = -\frac{1}{2} \sum_{\mu, \nu} \Gamma_{\mu\nu}^\alpha \frac{d\eta_\mu}{ds} \frac{d\eta_\nu}{ds}$$

gives the components of the gravitational field :

$$(15a) \quad \Gamma_{11}^1 = -\frac{1}{2} \frac{\partial f_1}{\partial \eta_1}$$

$$(15b) \quad \Gamma_{22}^1 = +\frac{1}{2} \frac{1}{f_1} \frac{\partial f_2}{\partial \eta_1} \frac{1}{1-\eta_2^2}$$

$$(15c) \quad \Gamma_{33}^1 = +\frac{1}{2} \frac{1}{f_1} \frac{\partial f_2}{\partial \eta_1} (1-\eta_2^2)$$

$$(15d) \quad \Gamma_{21}^2 = -\frac{1}{2} \frac{1}{f_2} \frac{\partial f_2}{\partial \eta_1}$$

$$(15e) \quad \Gamma_{22}^2 = -\frac{\eta_2}{1-\eta_2^2}$$

$$(15f) \quad \Gamma_{33}^2 = -\eta_2 (1-\eta_2^2)$$

$$(15g) \quad \Gamma_{31}^1 = -\frac{1}{2} \frac{1}{f_2} \frac{\partial f_2}{\partial \eta_1}$$

$$(15h) \quad \Gamma_{32}^2 = \frac{\eta_2}{1-\eta_2^2}$$

$$(15i) \quad \Gamma_{41}^4 = -\frac{1}{2} \frac{1}{f_4} \frac{\partial f_4}{\partial \eta_1}$$

The other ones are zero. Due to rotational symmetry it is sufficient to write the field equations only for the equator ($\eta_2=0$), therefore, since they will be differentiated only once, in the previous expressions it is possible to set everywhere since the beginning $1-\eta_2^2=0$. Then the calculation of the field equation gives :

$$(16a) \quad \frac{\partial}{\partial \eta_1} \left(\frac{1}{f_1} \frac{\partial f_1}{\partial \eta_1} \right) = \frac{1}{2} \left(\frac{1}{f_1} \frac{\partial f_1}{\partial \eta_1} \right)^2 + \left(\frac{1}{f_2} \frac{\partial f_2}{\partial \eta_1} \right)^2 + \left(\frac{1}{f_4} \frac{\partial f_4}{\partial \eta_1} \right)^2$$

$$(16b) \quad \frac{\partial}{\partial \eta_1} \left(\frac{1}{f_1} \frac{\partial f_2}{\partial \eta_1} \right) = 2 + \frac{1}{f_1 f_2} \left(\frac{\partial f_2}{\partial \eta_1} \right)^2$$

$$(16c) \quad \frac{\partial}{\partial \eta_1} \left(\frac{1}{f_1} \frac{\partial f_4}{\partial \eta_1} \right) = \frac{1}{f_1 f_4} \left(\frac{\partial f_4}{\partial \eta_1} \right)^2$$

Besides these three equations the functions f_1, f_2, f_3 must fullfill the equation of the determinant :

$$(17) \quad f_1 f_2^2 f_4 = 1 \quad i.e. \quad \frac{1}{f_1} \frac{\partial f_1}{\partial \eta_1} + \frac{2}{f_2} \frac{\partial f_2}{\partial \eta_1} + \frac{1}{f_4} \frac{\partial f_4}{\partial \eta_1} = 0$$

For now we neglect (16b) and determine the three functions f_1, f_2, f_4 from (16a), (16c) and (17). The equation (16c) can be transposed into the form :

$$(18) \quad \frac{\partial}{\partial \eta_1} \left(\frac{1}{f_4} \frac{\partial f_4}{\partial \eta_1} \right) = \frac{1}{f_1 f_4} \frac{\partial f_1}{\partial \eta_1} \frac{\partial f_4}{\partial \eta_1}$$

This can be integrated and gives

$$(19) \quad \frac{1}{f_4} \frac{\partial f_4}{\partial \eta_1} = \alpha f_1 \quad (\alpha \text{ being an integration constant})$$

The addition of (12a) and (12c') gives :

$$(20) \quad \frac{\partial}{\partial \eta_1} \left(\frac{1}{f_1} \frac{\partial f_1}{\partial \eta_1} + \frac{1}{f_4} \frac{\partial f_4}{\partial \eta_1} \right) = \left(\frac{1}{f_2} \frac{\partial f_2}{\partial \eta_1} \right)^2 + \frac{1}{2} \left(\frac{1}{f_1} \frac{\partial f_1}{\partial \eta_1} + \frac{1}{f_4} \frac{\partial f_4}{\partial \eta_1} \right)^2$$

By taking (17) into account we get :

$$(21) \quad -2 \frac{\partial}{\partial \eta_1} \left(\frac{1}{f_2} \frac{\partial f_2}{\partial \eta_1} \right) = 3 \left(\frac{1}{f_2} \frac{\partial f_2}{\partial \eta_1} \right)^2$$

By integrating :

$$(22) \quad \frac{1}{\frac{1}{f_2} \frac{\partial f_2}{\partial \eta_1}} = \frac{3}{2} \eta_1 + \frac{\sigma}{2} \quad (\sigma \text{ integration constant})$$

or :

$$(23) \quad \frac{1}{f_2} \frac{\partial f_2}{\partial \eta_1} = \frac{1}{3\eta_1 + \sigma}$$

After a second integration :

$$(24) \quad f_2 = \lambda (3\eta_1 + \sigma)^{2/3} \quad (\lambda \text{ integration constant})$$

The condition at infinity requires $\lambda = 1$. Then

$$(19) \quad f_2 = (3\eta_1 + \sigma)^{2/3}$$

Hence it results further from (22) and (17)

$$(20) \quad \frac{\partial f_4}{\partial \eta_1} = \alpha f_1 f_4 = \frac{\alpha}{f_2^2} = \frac{\alpha}{(3\eta_1 + \sigma)^{4/3}}$$

By integrating, while taking account the condition at infinity

$$(25) \quad f_4 = 1 - \alpha(3\eta_1 + \sigma)^{-1/3}$$

Hence, from (17)

$$(26) \quad f_1 = \frac{(3\eta_1 + \sigma)^{-4/3}}{1 - \alpha(3\eta_1 + \sigma)^{-1/3}}$$

As it can be easily verified the equation (16b) is automatically fulfilled by the expression that we found for f_1 and f_2

Therefore all the conditions are satisfied except the *condition of continuity*.

$$f_1 \text{ will be discontinuous when } 1 = \alpha(3\eta_1 + \alpha^3)^{-1/3}, \quad 3\eta_1 = \alpha^3 - \sigma$$

In order this discontinuity coincides with the origin, it must be :

$$(27) \quad \sigma = \alpha^3$$

Therefore the condition of continuity relates in this way the two integration constants σ and α . We have :

$$(28) \quad 3\eta_1 + \alpha = \zeta^3 + \alpha$$

$$(29) \quad f_4 = 1 - \frac{\alpha}{(\zeta^3 + \alpha^3)^{1/3}}$$

$$(30) \quad f_2 = (\zeta^3 + \alpha^3)^{2/3}$$

$$(31) \quad \frac{d\eta_2^2}{1 - \eta_2^2} = d\theta^2$$

Finally :

$$(32)$$

$$ds^2 = \left(1 - \frac{\alpha}{(\zeta^3 + \alpha^3)^{1/3}}\right) d\xi_0^2 - \frac{\zeta^4}{(\alpha^3 + \zeta^3) \left[(\alpha^3 + \zeta^3)^{1/3} - \alpha \right]} d\zeta^2 - (\alpha^3 + \zeta^3)^{2/3} (d\theta^2 + \sin^2 \theta d\phi^2)$$

When $\zeta \rightarrow \infty$ the line element tends to Lorentz form.

g_{tt} tends to zero when ζ tends to zero.

When $\zeta \rightarrow 0$ $g_{\zeta\zeta} \approx \frac{0}{0}$. An expansion into a series shows that $g_{\zeta\zeta} \approx \frac{3\zeta}{\alpha} \rightarrow 0$

Consider a loop located in the plane $\theta = 0$, with $d\xi_0 = 0$. It is no contractile : for $\zeta = 0$ the

+ jean-pierre.petit@manaty.net , gilles.dagostini@manaty.net, sebastien.michea@manaty.net

perimeter of the loop tends to $2\pi\alpha$.

Notice that $\zeta = \sqrt{\xi_1^2 + \xi_2^2 + \xi_3^2}$ is definitely not a « radius ». The point $\zeta = 0$ does not correspond to some « center of symmetry ». The spherically symmetric requirement does not identify automatically to a central symmetry, as suggested by David Hilbert [5] ¹ ζ is just one of the « space markers », nothing else. It's a number, not a length. The only length to be considered is the quantity s .

Expressing the metric in a better coordinate system.

Let us consider the coordinate system introduced in [3].

Introduce the new space marker ρ through the following coordinate change :

$$(33) \quad \zeta = |\alpha| \left(1 + \text{Logcosh } \rho \right)$$

$$(34) \quad \rho = \pm \text{argch} \left(e^{\sqrt[3]{1 + \left(\frac{\zeta}{|\alpha|}\right)^3} - 1} \right)$$

$$\zeta = 0 \quad \rightarrow \quad \rho = 0$$

With $\xi_0 = ct$ the line element becomes :

$$(35) \quad ds^2 = \frac{\text{Logcosh } \rho}{1 + \text{Logcosh } \rho} c^2 dt^2 - \frac{1 + \text{Logcosh } \rho}{\text{Logcosh } \rho} \alpha^2 \tanh^2 \rho d\rho^2 - \alpha^2 (1 + \text{Logcosh } \rho)^2 (d\theta^2 + \sin^2 \theta d\varphi^2)$$

When $\rho \rightarrow \pm \infty$ we get the following Lorentz form :

$$(36) \quad ds^2 = c^2 dt^2 - \alpha^2 d\rho^2 - \alpha^2 \rho^2 (d\theta^2 + \sin^2 \theta d\varphi^2)$$

We can figure this geometrical objet as a space bridge linking two Minkowski spacetimes, though a throat sphere whose perimeter is $2\pi\alpha$. We cannot think about its « radius » because that sphere has no center.

When $\rho \rightarrow 0$

¹ We quote, page 67 of the german edition : « Die Gravitation $g_{\mu\nu}$ ist zentrisch symmetrisch in Bezug auf den Kooridatenanfangspunkt. «

English translation : « The gravitation $g_{\mu\nu}$ is centrally symmetric with respect to the origin of coordinates. »

$$(37) \quad g_{tt} = \frac{\text{Log cosh } \rho}{1 + \text{Log cosh } \rho} c^2 \rightarrow 0$$

$$(38) \quad g_{\rho\rho} = -\frac{1 + \text{Log cosh } \rho}{\text{Log cosh } \rho} \alpha^2 \tanh^2 \rho \rightarrow \frac{0}{0}$$

$$(39) \quad g_{\theta\theta} \rightarrow \alpha^2 \quad g_{\varphi\varphi} \rightarrow \alpha^2 \sin^2 \theta$$

$$(40) \quad g_{\varphi\varphi} = \alpha^2 \sin^2 \theta$$

We may easily overcome the indetermination (33) through an expansion into a series, which shows that when $\rho \rightarrow 0$

$$(41) \quad g_{\rho\rho} = -\frac{1 + \text{Log cosh } \rho}{\text{Log cosh } \rho} \alpha^2 \tanh^2 \rho \rightarrow -2$$

The determinant is :

$$(42) \quad \det g_{\mu\nu} = -c^2 \alpha^6 \tanh^2 \rho \sin^2 \theta$$

It vanishes on the throat sphere. As a consequence, on this last we cannot define gaussian coordinates, so that the object is no longer a manifold but an orbifold. On the throat sphere the arrow of time and the space orientation cannot be defined. This can be interpreted as a geometric structure where space and time are reversed through the throat sphere : when a particle crosses the throat sphere it experiences a PT-symmetry. According to Souriau's theorem [4] this T-inversion goes with a mass inversion.

In a future paper the physical interpretation of such solution will be investigated.

References :

[1] K. Schwarzschild : Über das Gravitationsfeld einer Kugel Aus incompressibler Flüssigkeit nach der Einsteinschen Theorie. Sitzung der phys. Math. Klasse v.23 märz 1916

[2] K. Schwarzschild : On the gravitational field of a sphere of incompressible fluid according to Einstein theory. Translation by S.Antoci and A.Loinger. arXiv :physics/9905030v1 [physics.hist-ph] 12 may 1999.

[3] J.P.Petit & G.D'Agostini : Cancellation of the singularity of the Schwarzschild solution with natural mass inversion process. Mod. Phys. Lett. A vol. 30 n°9 2015

[4] J.M.Souriau : Structure des systèmes dynamiques. Dunod Ed. France, 1970 and Structure of Dynamical Systems. Boston, Birkhäuser Ed. 1997. For time inversion see page 190 equation (14.67).

[5] D.Hilbert : “Die Grundlagen der Physij. (Zweite Mitteilung)” in Nachrichten von der Königlichen Gesellschaft zu Göttingen. Math. -Phys. Klasse. 1917 , p.53-76. Presented in the session of 23 december 1916